

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
20 January 2000 (20.01.2000)

PCT

(10) International Publication Number
WO 00/02522 A3(51) International Patent Classification: C12N 15/31,
C07K 14/32, A61K 39/07, C12N 7/01, 1/216936 Eylers Valley Flint Road, Sabillasville, MD 21780
(US). WELKOS, Susan, L. [US/US]; 1594 Dockside
Drive, Frederick, MD 21701 (US).

(21) International Application Number: PCT/US99/15568

(22) International Filing Date: 9 July 1999 (09.07.1999)

(74) Agents: HARRIS, Charles, II.; United States Army Medical
Research and Materiel Command, 504 Scott Street,
Fort Detrick, MD 21702 et al. (US).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60092,416 10 July 1998 (10.07.1998) US(71) Applicant (for all designated States except US): U.S.
MEDICAL RESEARCH INSTITUTE OF INFEC-
TIOUS DISEASES [US/US]; Department of the Army,
504 Scott Street, Frederick, MD 21702 (US).(81) Designated States (national): AL, AM, AT, AU, AZ, BA,
BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES,
FI, GB, GE, GH, GM, HU, ID, IL, IS, JP, KE, KG, KP, KR,
KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW,
MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,
TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW.(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM,
AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT,
BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,
NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA,
GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and


(75) Inventors/Applicants (for US only): LEE, John, S.
[US/US]; 1807 Elizabeth Court, Hagerstown, MD 21740
(US). PUSHKO, Peter [US/US]; 917 Seminole Road,
Frederick, MD 21701 (US). PARKER, Michael, D.
[US/US]; Apartment T-13, 1001 Carroll Parkway, Fred-
erick, MD 21701 (US). SMITH, Jonathan, F. [US/US];

Published:

— With international search report.

[Continued on next page]

(54) Title: ANTHRAX VACCINE



(57) Abstract: Using the nontoxic protective antigen PA protein from B. anthracis, a method and composition for use in inducing an immune response which is protective against anthrax in subjects is described.



(88) Date of publication of the international search report:
12 October 2000

(15) Information about Correction:
see PCT Gazette No. 14/2001 of 5 April 2001, Section II

(48) Date of publication of this corrected version:
5 April 2001

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

TITLE OF THE INVENTION

Anthrax Vaccine

5

10

FIELD OF THE INVENTION

This invention relates to vaccines for bacterial toxins from *Bacillus anthracis*.

15

INTRODUCTION

Anthrax is a disease cause by the sporulating bacteria *Bacillus anthracis*. Humans working with animal products are at risk from contracting anthrax. Areas such as Iran, Turkey, Iraq, Pakistan, and sub-Saharan Africa are hyperendemic for anthrax, although the organism can be found in most areas of the world. Anthrax manifests disease in three different ways. Inhalation, gastrointestinal, and cutaneous anthrax result from inhaling spores, ingesting spores in contaminated meat, or contacting spores in an open wound, respectively. Untreated inhalation or gastrointestinal anthrax has a case fatality rate of essentially 100 percent while cutaneous anthrax has a case fatality rate of up to 25 percent. Previous research has shown that the protective antigen (PA) produced by *B. anthracis* can protect mice from anthrax. Even though the Anthrax vaccine is FDA licensed, reactogenicity is mild to moderate.

Therefore, there is a need for an efficacious vaccine for anthrax useful for protecting humans.

SUMMARY OF THE INVENTION

5 The present invention satisfies the need discussed above. The present invention relates to a method and composition for use in inducing an immune response which is protective against infection with anthrax.

10 In this vaccine strategy, a gene coding for a protein of interest is cloned into a VEE virus vector in place of the VEE virus structural genes; the result is a self-replicating RNA molecule, a replicon, that encodes its own replicase and transcriptase functions,
15 and in addition makes abundant quantities of mRNA encoding the foreign protein. When replicon RNA is transfected into eukaryotic cells along with two helper RNAs that express the VEE structural proteins (glycoproteins and nucleocapsid), the replicon RNA is
20 packaged into VEE virus-like particles by the VEE virus structural proteins, which are provided in trans. Since the helper RNAs lack packaging signals necessary for further propagation, the resulting VEE replicon particles (VRPs) which are produced are
25 infectious for one cycle but are defective thereafter. Upon infection of an individual cell with a VRP, an abortive infection occurs in which the infected cell produces the protein of interest in abundance, is ultimately killed by the infection, but does not
30 produce any viral progeny (Pushko *et al.*, 1997, *Virology* 239, 389-401).

 The PA gene which contains a prokaryotic secretory signal and the entire 83 kDa coding sequence (Welkos *et al.*, 1988, *Gene* 69, 287-300), was inserted

into the VEE replicon vaccine vector (Figure 1) and have demonstrated high level expression of this bacterial protein in eukaryotic cells in culture. Mice, either the C57BL/6 strain or the A/J strain, inoculated with VRP containing the PA-replicon produced high specific antibody titers and were protected from developing anthrax when challenged subcutaneously with *B. anthracis*.

Therefore, it is one object of the present invention to provide a VEE virus replicon vector comprising a VEE virus replicon and DNA fragments encoding the PA protein from *B. anthracis*.

It is another object of the present invention to provide a self replicating RNA comprising the VEE virus replicon and any of the *B. anthracis* fragments described above.

It is another object of the present invention to provide infectious VEE virus replicon particles produced from the VEE virus replicon RNA described above.

It is further an object of the invention to provide an immunological composition for the protection of mammals against *B. anthracis* infection comprising VEE virus replicon particles containing any of the *B. anthracis* fragments described above or a combination of different VEE virus replicons each having a different *B. anthracis* fragment.

BRIEF DESCRIPTION OF THE DRAWINGS

These and other features, aspects, and advantages of the present invention will become better understood with reference to the following description and appended claims, and accompanying drawings where:

Figure 1. Diagram of the protective antigen (PA) replicon constructs. The PA replicons are similar to

the full-length VEE RNA except that the open reading frame encoding the VEE structural proteins was replaced with the PA genes.

Figure 2. Western blot of BHK cell lysates showing expression of anthrax protective antigen from recombinant VEE replicons. a) uninfected cell lysate; b) recombinant expression product from *B. anthracis*; c) infected cell lysate.

Figure 3A and 3B. ELISA titers for mice (A, C57Bl/6 mice; B, A/J mice) immunized with PA-VRP. GMTs only includes those mice that survived challenge. a) Received 0.2 ml on day 0 and 28. b) Received 10^7 iu on day 0 and 28. c) Received 10^7 iu PA-VRP on day 0, 28, 56, and 84; C57Bl/6 mice that died, titer = 100; A/J mice that died, titer = 400. d) Received 10^7 iu PA-VRP on day 0, 28, and 56; C57Bl/6 mice that died, titers = 400, 100, 100; A/J mice that died, titers = 102400, 100, 1600. e) Received 10^7 iu PA-VRP on day 0, and 28; C57Bl/6 mice that died, GMT = 1313; A/J mice that died, GMT = 1766. Mice were challenged 28 days after last inoculation with 18-32 LD₅₀ Sterne strain for C57Bl/6 and 12-25 LD₅₀ Sterne strain for A/J mice.

Figure 4. Schematic representation of constructs containig PA.

DETAILED DESCRIPTION

In the description that follows, a number of terms used in recombinant DNA, virology and immunology are extensively utilized. In order to provide a clearer and consistent understanding of the specification and claims, including the scope to be given such terms, the following definitions are provided.

Replicon. A replicon is equivalent to a full length virus from which all of the viral structural proteins have been deleted. A multiple cloning site can be cloned into the site previously occupied by the structural protein genes. Virtually any heterologous gene may be cloned into this cloning site. Transcription of the RNA from the replicon yields an RNA capable of initiating infection of the cell identically to that seen with the full-length infectious virus clone. However, in lieu of the viral structural proteins, the heterologous antigen is expressed. This system does not yield any progeny virus particles because there are no viral structural proteins available to package the RNA into particles.

Particles which appear structurally identical to virus particles can be produced by supplying structural proteins for packaging of the replicon RNA *in trans*. This is typically done with two helpers also called defective helper RNAs. One helper consists of a full length infectious clone from which the nonstructural protein genes and the glycoprotein genes are deleted. The helper retains only the terminal nucleotide sequences, the promoter for subgenomic mRNA transcription and the sequences for the viral nucleocapsid protein. The second helper is identical to the first except that the nucleocapsid gene is deleted and only the glycoprotein genes are retained. The helper RNA's are transcribed *in vitro* and co-transfected with replicon RNA. Because the replicon RNA retains the sequences for packaging by the nucleocapsid protein, and because the helpers lack these sequences, only the replicon RNA is packaged by the viral structural proteins and released from the cell. The particles can then be inoculated into animals similar to parent virus. The replicon

particles will initiate only a single round of replication because the helpers are absent, they produce no progeny virus particles, and express only the viral nonstructural proteins and the product of the heterologous gene cloned in place of the structural proteins.

The VEE virus replicon is a genetically reorganized version of the VEE virus genome in which the structural proteins genes are replaced with a gene from an immunogen of interest, in this invention, the *B. anthracis* proteins. The result is a self replicating RNA (replicon) that can be packaged into infectious particles using defective helper RNAs that encode the glycoprotein and capsid proteins of the VEE virus.

Subject. Includes both human, animal, e.g., horse, cattle, donkey, monkey, pig, dog, guinea pig, mouse, hamster, avian e.g., chicken, pheasant or turkey, fish and other marine animals, and insects such as mosquito.

In one embodiment, the present invention relates to a recombinant DNA molecule that includes a VEE replicon and a DNA sequence encoding the non-toxic protective antigen (PA) from *B. anthracis*. The sequence for PA has been determined and has been deposited in GenBank at accession no. M22589. The PA sequence encodes a prokaryotic secretory signal in addition to the entire sequence encoding the 83 Kd PA (SEQ ID NO:1). Other nucleic acid sequences related to this invention include sequences wherein the secretory signal has been removed (MAT-PA) (SEQ ID NO:2), or replaced with other secretory signals known to people in the art such as the tissue plasminogen activator (TPA) secretory signal resulting in a DNA

fragment encoding TPA-PA (SEQ ID NO:3). Nucleic acid sequences included in this invention are sequences encoding the active form of PA, a 63 kDa protein, termed PA63 (SEQ ID NO:4). In addition, isolated
5 nucleic acid molecules of the invention include DNA molecules which comprise a sequence substantially different from those described above but which, due to the degeneracy of the genetic code, still encode the *B. anthracis* proteins described and specified in
10 SEQ ID NO:5 (TPA-PA), SEQ ID NO:6 (PA), SEQ ID NO:7 (MAT-PA), and SEQ ID NO:8 (PA63). Of course, the genetic code and species-specific codon preferences are well known in the art. Thus, it would be routine for one skilled in the art to generate the degenerate
15 variants described above, for instance, to optimize codon expression for a particular host (e.g., change codons in the human mRNA to those preferred by a bacterial host such as *E.coli*).

Nucleic acid molecules of the present invention
20 may be in the form of RNA, or in the form of DNA, including, for instance, cDNA and genomic DNA obtained by cloning or produced synthetically. The DNA may be double-stranded or single-stranded. Single-stranded DNA or RNA may be the coding strand, also known as the
25 sense strand, or it may be the non-coding strand, also referred to as the antisense strand.

By "isolated" nucleic acid molecule(s) is intended a nucleic acid molecule, DNA or RNA, which has been removed from its native environment. For
30 example, recombinant DNA molecules contained in a vector are considered isolated for the purposes of the present invention. Further examples of isolated DNA molecules include recombinant DNA molecules maintained in heterologous host cells or purified (partially or
35 substantially) DNA molecules in solution. Isolated

RNA molecules include *in vivo* or *in vitro* RNA transcripts of the DNA molecules of the present invention. Isolated nucleic acid molecules according to the present invention further include such molecules produced synthetically.

The present invention is further directed to nucleic acid molecules encoding portions or fragments of the nucleotide sequences described herein. Fragments include portions of the nucleotide sequences of at least 10 contiguous nucleotides in length selected from any two integers, one of which representing a 5' nucleotide position and a second of which representing a 3' nucleotide position, where the first nucleotide for each nucleotide sequence is position 1. That is, every combination of a 5' and 3' nucleotide position that a fragment at least 10 contiguous nucleotide bases in length or any integer between 10 and the length of an entire nucleotide sequence minus 1.

The present invention further relates to variants of the nucleic acid molecules of the present invention, which encode portions, analogs or derivatives of the *B. anthracis* polypeptides described above. Variants may occur naturally, such as a natural allelic variant. By an "allelic variant" is intended one of several alternate forms of a gene occupying a given locus of a chromosome of an organism. Non-naturally occurring variants may be produced by known mutagenesis techniques. Such variants include those produced by nucleotide substitution, deletion, or addition of one or more nucleotides in the coding or noncoding regions or both. Alterations in the coding regions may produce conservative or nonconservative amino acid substitutions, deletions, or additions. Especially

preferred among these are silent substitutions, additions, and deletions which do not alter the properties and activities of the *B. anthracis* polypeptides disclosed herein or portions thereof.

- 5 Also preferred in this regard are conservative substitutions.

In another embodiment, the present invention relates to a recombinant DNA molecule that includes a vector and a DNA sequence as described above. The
10 vector can take the form of a plasmid, phage, cosmid, YAC, eukaryotic expression vector such as a DNA vector, *Pichia pastoris*, or a virus vector such as for example, baculovirus vectors, retroviral vectors or adenoviral vectors, and others known in the art. The
15 cloned gene may optionally be placed under the control of (i.e., operably linked to) certain control sequences such as promoter sequences, or sequences which may be inducible and/or cell type-specific. Suitable promoters will be known to a person with
20 ordinary skill in the art. The expression construct will further contain sites for transcription initiation, termination and, in the transcribed region, a ribosome binding site for translation. When the DNA sequences described above are in a
25 replicon expression system, such as the VEE replicon described above, the proteins can be expressed *in vivo*. The DNA sequence for any of the *B. anthracis* proteins described above can be cloned into the multiple cloning site of a replicon such that
30 transcription of the RNA from the replicon yields an infectious RNA containing the *B. anthracis* protein or proteins of interest. Use of helper RNA containing sequences necessary for encapsulation of the viral transcript will result in the production of viral
35 particles containing replicon RNA which are able to

infect a host and initiate a single round of replication resulting in the expression of the *B. anthracis* proteins. Such replicon constructs include those listed in Table 1.

5 Table 1

Replicon	expresses
<u>Plasmid Name</u>	
p3014-TPA-PA	TPA signal, 5 amino acids of C terminal end of prokaryotic secretory signal, and the 83kDa mature protein.
p3014-PA	29 amino acid prokaryotic secretory signal, and the 83kDa mature protein secreted by <i>B. anthracis</i>
p3014-MAT-PA	83kDa mature protein
p3014-PA63	63kDa protein (minus 20kDa N-terminal end that is naturally cleaved off by furin proteases at host cell membrane surface)
p3014-sub-domains	subdomain #1: from nucleotide 1891 to 2391; subdomain #2: from nucleotide 2392 to 2658; subdomain #3: from nucleotide 2659 to 3351; subdomain #4: from nucleotide 3352 to 3669; subdomain #5: from nucleotide 3670 to 4098 (Subdomains are numbered according to Welkos et al., 1988, supra, determined on subdomains present in the crystal structure, Petosa, et al., 1997, Nature 385, 833-838).

The VEE constructs containing anthrax proteins can be used as a DNA vaccine, or for the production of RNA molecules as described below.

40 In another embodiment, the present invention relates to RNA molecules resulting from the transcription of the constructs described above. The RNA molecules can be prepared by *in vitro* transcription using methods known in the art and
 45 described in the Examples below. Alternatively, the

RNA molecules can be produced by transcription of the constructs *in vivo*, and isolating the RNA. These and other methods for obtaining RNA transcripts of the constructs are known in the art. Please see Current
5 Protocols in Molecular Biology. Frederick M. Ausubel et al. (eds.), John Wiley and Sons, Inc. The RNA molecules can be used, for example, as a nucleic acid vaccine, or to transfect cells along with RNA from helper plasmids, one of which expresses VEE
10 glycoproteins and the other VEE capsid proteins, as described above, in order to obtain replicon particles.

Introduction of the construct into the host cell can be effected by calcium phosphate transfection,
15 electroporation, infection, and other methods known in the art and described in standard laboratory manuals such as Current Protocols in Molecular Biology, Ausubel, F. M. et al. (Eds), Wiley & Sons, Inc. All documents cited herein supra and infra are hereby
20 incorporated in their entirety by reference thereto.

In a further embodiment, the present invention relates to host cells stably transformed or transfected with the above-described recombinant DNA constructs. The host cell can be prokaryotic (for
25 example, bacterial), lower eukaryotic (for example, yeast or insect) or higher eukaryotic (for example, all mammals, including but not limited to rat and human). Both prokaryotic and eukaryotic host cells may be used for expression of desired coding sequences
30 when appropriate control sequences which are compatible with the designated host are used. Among prokaryotic hosts, *E. coli* is most frequently used. Expression control sequences for prokaryotes include promoters, optionally containing operator portions,

and ribosome binding sites. Transfer vectors compatible with prokaryotic hosts are commonly derived from, for example, pBR322, a plasmid containing operons conferring ampicillin and tetracycline resistance, and the various pUC vectors, which also contain sequences conferring antibiotic resistance markers. These markers may be used to obtain successful transformants by selection. Please see e.g., Maniatis, Fritsch and Sambrook, Molecular Cloning: A Laboratory Manual (1982) or DNA Cloning, Volumes I and II (D. N. Glover ed. 1985) for general cloning methods. The DNA sequence can be present in the vector operably linked to a sequence encoding an IgG molecule, an adjuvant, a carrier, or an agent for aid in purification of protein of the invention, such as glutathione S-transferase. The recombinant molecule can be suitable for transfecting eukaryotic cells, for example, mammalian cells and yeast cells in culture systems. *Saccharomyces cerevisiae*, *Saccharomyces carlsbergensis*, and *Pichia pastoris* are the most commonly used yeast hosts, and are convenient fungal hosts. Control sequences for yeast vectors are known in the art. Mammalian cell lines available as hosts for expression are known in the art and include many immortalized cell lines available from the American Type Culture Collection (ATCC), such as baby hamster kidney (BHK) cells, MRC-5 cells and vero cells, to name a few. Suitable promoters are also known in the art and include viral promoters such as that from SV40, Rous sarcoma virus (RSV), adenovirus (ADV), bovine papilloma virus (BPV), and cytomegalovirus (CMV). Mammalian cells may also require terminator sequences and poly A addition sequences; enhancer sequences which increase

expression may also be included, and sequences which cause amplification of the gene may also be desirable. These sequences are known in the art. The transformed or transfected host cells can be used as a source of DNA sequences described above. When the recombinant molecule takes the form of an expression system, the transformed or transfected cells can be used as a source of the protein or polypeptide specified as amino acid sequence in SEQ ID NO:5 (TPA-PA), SEQ ID NO:6 (PA), SEQ ID NO:7 (MAT-PA), and SEQ ID NO:8 (PA63).

A polypeptide or amino acid sequence derived from the amino acid sequences mentioned above, refers to a polypeptide having an amino acid sequence identical to that of a polypeptide encoded in the sequence, or a portion thereof wherein the portion consists of at least 2-5 amino acids, and more preferably at least 8-10 amino acids, and even more preferably at least 11-15 amino acids, or which is immunologically identifiable with a polypeptide encoded in the sequence.

A recombinant or derived polypeptide is not necessarily translated from a designated nucleic acid sequence; it may be generated in any manner, including for example, chemical synthesis, or expression of a recombinant expression system. In addition the polypeptide can be fused to other proteins or polypeptides which increase its antigenicity, such as adjuvants for example.

The recombinant or fusion protein can be used as a vaccine for immunity against anthrax infection or as a diagnostic tool for detection of bacillus anthracis. The transformed host cells can be used to analyze the effectiveness of drugs and agents which inhibit anthrax or *B. anthracis* proteins, such as host

proteins or chemically derived agents or other proteins which may interact with *B. anthracis* proteins of the present invention to inhibit its function. A method for testing the effectiveness of an anti-anthrax drug or agent can for example be the rat anthrax toxin assay (Ivins et al. 1984, *Infect. Immun.* 52, 454-458 and Ivins et al. 1986) or a skin test in rabbits for assaying antiserum against anthrax toxin (Belton and Henderson, 1956, *Br. J. Exp. Path.* 37, 156-160).

In another embodiment, the present invention relates to an anthrax vaccine comprising one or more replicon particles derived from one or more replicons encoding one or more *B. anthracis* proteins or polypeptides as described above. The present invention relates a method for providing immunity against anthrax said method comprising administering one or more replicon particles containing any combination of the *B. anthracis* proteins to a subject such that a protective immune reaction is generated.

Vaccine formulations of the present invention comprise an immunogenic amount of a replicon particle, resulting from one of the replicon constructs described above, or a combination of replicon particles as a multivalent vaccine, in combination with a pharmaceutically acceptable carrier. An "immunogenic amount" is an amount of the replicon particles sufficient to evoke an immune response in the subject to which the vaccine is administered. An amount of from about 10^2 to 10^7 per dose is suitable, more or less can be used depending upon the age and species of the subject being treated. Exemplary pharmaceutically acceptable carriers include, but are not limited to, sterile pyrogen-free water and sterile pyrogen-free physiological saline solution.

Administration of the replicon particles disclosed herein may be carried out by any suitable means, including both parenteral injection (such as intraperitoneal, subcutaneous, or intramuscular injection), by in ovo injection in birds, orally and by topical application of the virus (typically carried in the pharmaceutical formulation) to an airway surface. Topical application of the virus to an airway surface can be carried out by intranasal administration (e.g. by use of dropper, swab, or inhaler which deposits a pharmaceutical formulation intranasally). Topical application of the virus to an airway surface can also be carried out by inhalation administration, such as by creating respirable particles of a pharmaceutical formulation (including both solid particles and liquid particles) containing the replicon as an aerosol suspension, and then causing the subject to inhale the respirable particles. Methods and apparatus for administering respirable particles of pharmaceutical formulations are well known, and any conventional technique can be employed. An "immunogenic amount" is an amount of the replicon particles sufficient to evoke an immune response in the subject to which the vaccine is administered.

When the replicon RNA or DNA is used as a vaccine, the replicon RNA or DNA can be administered directly using techniques such as delivery on gold beads (gene gun), delivery by liposomes, or direct injection, among other methods known to people in the art. Any one or more constructs or replicating RNA described above can be use in any combination effective to elicit an immunogenic response in a subject. Generally, the nucleic acid vaccine administered may be in an amount of about 1-5 ug of

nucleic acid per dose and will depend on the subject to be treated, capacity of the subject's immune system to develop the desired immune response, and the degree of protection desired. Precise amounts of the vaccine
5 to be administered may depend on the judgement of the practitioner and may be peculiar to each subject and antigen.

The vaccine may be given in a single dose schedule, or preferably a multiple dose schedule in
10 which a primary course of vaccination may be with 1-10 separate doses, followed by other doses given at subsequent time intervals required to maintain and or reinforce the immune response, for example, at 1-4 months for a second dose, and if needed, a subsequent
15 dose(s) after several months. Examples of suitable immunization schedules include: (i) 0, 1 months and 6 months, (ii) 0, 7 days and 1 month, (iii) 0 and 1 month, (iv) 0 and 6 months, or other schedules
20 sufficient to elicit the desired immune responses expected to confer protective immunity, or reduce disease symptoms, or reduce severity of disease.

The following MATERIALS AND METHODS were used in the examples that follow.

Plasmids. Construction of the VEE replicon, capsid (C-) helper, and glycoprotein (GP-) helper
25 plasmids was previously described (Pushko, 1997, supra). The PA gene from nucleotide 1804 to 4098 (SEQ ID NO:1) (Welkos et al., 1988, supra) was cloned into the VEE replicon plasmid as a BamHI/BamHI fragment
30 utilizing a shuttle vector. TPA-PA (SEQ ID NO:3), MAT-PA (SEQ ID NO:2), and PA63 (SEQ ID NO:4) were PCR amplified using forward and reverse primers which also contained the ClaI recognition site. The PCR products were agarose gel purified and ligated into the T/A
35 vector pCR2.1 (Invitrogen, Inc.). The PA genes, after

digestion of pCR2.1-PA's with ClaI, were ligated into the VEE replicon plasmid. The Lassa nucleocapsid replicon (Lassa N-replicon) was constructed as previously described (Pushko, 1997, supra) and used as a negative control replicon.

Production of VRP. Plasmid templates for the PA-replicons, C-helper, GP-helper, and the Lassa N-replicon were linearized by digestion with NotI at a unique site downstream from the coding sequences, and capped run-off transcripts were prepared *in vitro* using T7 RNA polymerase. Packaging of the replicons into VEE replicon particles (VRPs) was accomplished by electroporating the replicon RNA and the two helper RNAs into BHK cells. VRPs were harvested between 20 and 27 hours after transfection and purified from cell culture supernatants by ultracentrifugation through a discontinuous sucrose gradient (20%). After reconstituting the pelleted VRP in 1/50 volume phosphate buffered saline, the VRPs were stored at -70°C.

Analysis of expression products and titration of VRP. Subconfluent monolayers were infected with PA-VRP's (either TPA-PA, PA, MAT-PA, or PA63) or Lassa N-VRP (m.o.i. = 2) or cell suspensions were electroporated with replicon RNA. Cells were harvested at approximately 20-24 hours and expressed proteins were separated by SDS-PAGE. Visualization of PA proteins was accomplished using a chemiluminescence western blot assay and antibodies specific for each protein. Titration of VRPs was accomplished by infecting subconfluent monolayers with increasing dilutions of purified VRP. Antigen positive cells were visualized in an indirect immunofluorescence assay using a monoclonal antibody specific for each protein,

or in a direct immunofluorescence assay using an FITC-conjugated monkey anti-Lassa serum.

Immunization of mice. Mice were inoculated 1 to 4 times at 28 day intervals with 10^5 to 10^7 infectious units (iu) of either PA-VRP or Lassa N-VRP (negative control). Positive control mice for the anthrax study were inoculated subcutaneously with 0.2 ml of anthrax vaccine at 28 day intervals. Serum for ELISA was obtained 2 days before each inoculation and 3 days before challenge.

Enzyme-linked immunosorbent assay (ELISA). Microtiter plates were coated with protective antigen (gift from Joe Farchaus, Bacteriology Division) (1 ug/ml) in PBS and allowed to absorb overnight at 4°C. Four fold serum dilutions in blocking buffer were applied to the plates and incubated at 37°C for 1 hour. After washing, an anti-mouse secondary antibody (HRP conjugated) was added to the plate and incubated for an additional hour at 37°C. After washing, bound antibody was detected colorimetrically using ABTS as a substrate.

Challenge of mice.

Anthrax challenge: C57BL/6 or A/J mice were challenged subcutaneously with 12-32 LD₅₀ *B. anthracis* (Sterne) 31 days after the last inoculation.

Example 1

Packaging and expression of PA-replicon

The PA-replicon was efficiently packaged into VRPs using the double helper system. Stock solutions contained about 10^8 infectious units of purified PA-VRP per milliliter. No replication competent VEE virus was detected in any of the preparations. A western blot of cell lysates, generated from BHK cells infected with PA-VRP, demonstrates a product of

approximately 85 kDa (Figure 2). Because the PA-VRP expression product contains a 29 amino acid prokaryotic secretory signal, which is cleaved from the standard, the PA product from the BHK cells appears larger than the PA standard.

Example 2

Protection against challenge with *B. anthracis*

The results of the animal studies demonstrated that PA from *B. anthracis* could be used to immunize and protect mice from a lethal challenge of *B. anthracis* (Sterne strain). Although the Sterne strain is attenuated in some animals, it is virulent in mice. Controls in the experiment included mice vaccinated with the current human vaccine (anthrax vaccine absorbed, AVA) which protected all mice, and mice immunized with the Lassa N-replicon which expressed an antigen irrelevant to anthrax protection which failed to protect the mice. Figure 3A shows ELISA titers and survival for C57Bl/6 mice inoculated with two to four doses of PA-VRP. Seventy percent of the C57Bl/6 mice which received three inoculations of the PA-VRP were protected whereas 90% of the C57Bl/6 mice which received four inoculations were protected from an otherwise lethal challenge of *B. anthracis* (Sterne strain). The geometric mean of the serum ELISA titers to PA of C57Bl/6 mice given four doses of the PA-replicon was 150500, as compared to 264 for mice that were inoculated with the negative control replicon. Figure 3B shows ELISA titers and survival for A/J mice inoculated with two to four doses of PA-VRP. Three inoculations of the PA-VRP protected 70% of the A/J mice, whereas four inoculations protected 90% of the A/J mice from an otherwise lethal challenge of the *B. anthracis* (Sterne strain). The geometric mean of the serum ELISA titers against PA in A/J mice given four

doses of the PA-replicon was 223336 as compared to 100 for mice which were inoculated with the negative control replicon. As expected, none of the mice (from either strain) which were inoculated with the negative control replicon survived challenge with *B. anthracis* (Sterne strain).

Discussion/Conclusion

Since VEE virus replicates in the cytoplasm of eukaryotic cells, the VEE replicon vaccine vector is a useful tool for the expression of prokaryotic genes in eukaryotic cells. Cytoplasmic expression of genes alleviates the difficulties imposed by splicing and nuclear transport of mRNA. We used the VEE replicon as a way to express the prokaryotic PA gene in eukaryotic cells and to develop a new vaccine candidate against anthrax.

The human AVA contains mostly PA protein and is presumed to protect humans by eliciting an antibody response that can neutralize the PA portion of anthrax toxin produced by invading *B. anthracis*. After neutralization of the toxin, the immune system destroys the invading bacteria. In order to model the human disease, we chose to use mice and the Sterne strain of *B. anthracis*. The Sterne strain produces anthrax toxin (encoded on the pX01 plasmid) but lacks a capsule (encoded on the pX02 plasmid). Mice exposed to large quantities of Sterne (ranging from 10^3 to 10^8 spores for the different mouse strains) are not able to overcome the effects of the toxin produced by the bacteria so they succumb to infection. We hypothesized that by immunizing the mice with PA, thereby inducing neutralizing antibodies to the PA

portion of the toxin produced by a challenge inoculum, that the mice would be protected.

A VEE replicon was constructed which expressed the PA gene from *B. anthracis* and was evaluated for *in vitro* expression, *in vivo* immunogenicity, and protective efficacy. C57Bl/6 and A/J mice demonstrated increasing antibody responses to PA when inoculated with increasing doses of PA-VRP. The two mouse strains inoculated with up to four doses of PA-VRP displayed increased protection against a lethal challenge with *B. anthracis* (Sterne strain). The use of mice and the Sterne strain of *B. anthracis* useful as a model system for studying the immunogenicity and efficacy of replicon based anthrax vaccines.

15

20

25

30

What is claimed is:

1. A recombinant DNA construct comprising:
 - (i) a vector, and
 - (ii) at least one nucleic acid
- 5 fragment comprising any combination of *B. anthracis* proteins selected from the group consisting of PA, MAT-PA, TPA-PA, and PA63.
2. A recombinant DNA construct according to claim
- 10 1 wherein said vector is an expression vector.
3. A recombinant DNA construct according to claim
- 1 wherein said vector is a prokaryotic vector.
- 15 4. A recombinant DNA construct according to claim
- 1 wherein said vector is a eukaryotic vector.
5. The recombinant DNA construct of claim 1
- wherein said vector is a VEE virus replicon vector.
- 20 6 The recombinant DNA construct according to
- claim 5 wherein said construct is p3014-PA.
7. The recombinant DNA construct according to
- 25 claim 5 wherein said construct is p3014-TPA-PA.
8. The recombinant DNA construct according to
- claim 5 wherein said construct is p3014-MAT-PA.
- 30 9. The recombinant DNA construct according to
- claim 5 wherein said construct is p3014-PA63.
10. Self replicating RNA produced from a
- construct chosen from the group consisting of: p3014-
- 35 PA, p3014-TPA-PA, p3014-MAT-PA, and p3014-P63

11. Infectious alphavirus particles produced from packaging the self replicating RNA of claim 10.
- 5 12. A pharmaceutical composition comprising infectious alphavirus particles according to claim 11 in an effective immunogenic amount in a pharmaceutically acceptable carrier and/or adjuvant.
- 10 13. A host cell transformed with a recombinant DNA construct according to claim 5.
14. A host cell according to claim 13 wherein said host cell is prokaryotic.
- 15 15. A host cell according to claim 13 wherein said host cell is eukaryotic.
16. A method for producing *B. anthracis* protein comprising culturing the cells according to claim 14 under conditions such that said DNA fragment is expressed and said protein is produced.
- 20 17. A method for producing *B. anthracis* protein comprising culturing the cells according to claim 15 under conditions such that said DNA fragment is expressed and said protein is produced.
- 25 18. A vaccine for anthrax comprising viral particles containing one or more replicon RNA encoding any combination of *B. anthracis* proteins chosen from the group consisting of PA, PA-MAT, TPA-PA, and PA63.
- 30 19. A pharmaceutical composition comprising the self replication RNA of claim 14 in an effective
- 35

immunogenic amount in a pharmaceutically acceptable carrier and/or adjuvant.

20. A pharmaceutical composition comprising one
5 or more recombinant DNA constructs chosen from the
group consisting of p3014-PA, p3014-TPA-PA, p3014-MAT-
PA, and p3014-P63 in a pharmaceutically acceptable
amount, in a pharmaceutically acceptable carrier/and
or adjuvant.

10

15

20

25

30

FIG. 1




FIG. 2

lysate ^a	STD ^b	lysate ^c	lysate ^c	lysate ^c	Secretory Sequence
	-	+	+	+	



FIG. 3A




FIG. 3B





FIG. 4



SEQUENCE LISTING

<110> United States Army Medical Research Institute of
Infectious Diseases
Lee, John Scott
Pushko, Peter
Parker, Michael D.
Smith, Jonathan F.
Welkos, Susan L.

<120> Anthrax Vaccines

<130> 003/135/SAP

<140> US 09/350,729

<141> 1999-07-09

<150> US 60/092,416

<151> 1998-07-10

<160> 8

<170> Apple Macintosh Microsoft Word 6.0

<210> 1

<211> 2295

<212> DNA

<213> *Bacillus anthracis*

<220>

<400> 1

atgaaaaaac	gaaaagtgtt	aataccatta	atggcattgt	40
ctacgatatt	agtttcaagc	acaggtaatt	tagaggat	80
tcaggcagaa	gttaaacagg	agaacccgtt	attaaatgaa	120
tcagaatcaa	gttccagggt	gttactagga	tactatttta	160
gtgatttgaa	ttttcaagca	cccattgttg	ttacctcttc	200
tactacaggg	gatttatcta	ttcctagtct	tgagttagaa	240
aattattccat	cggaaaacca	atattttcaa	tctgctattt	280
ggtcaggatt	tatcaaaagt	aagaagagtg	atgaatatatc	320
atttgctact	tccgctgata	atcatgtaac	aatgtgggta	360
gatgaccaag	aagtgatata	taaagcttct	aatttcaaca	400
aaatcagatt	agaaaaagga	agattatatc	aaataaaaat	440
tcaatatcaa	cgagaaaatc	ctactgaaaa	aggattggat	480
ttcaagtgtg	actggaccga	ttctcaaaat	aaaaaagaag	520
tgattttctag	tgataactta	caattgccag	aattaaaaca	560
aaaattcttcg	aactcaagaa	aaaagacaag	tacaagtgc	600


```

ggacctacgg  ttccagaccg  tgacaatgat  ggaatccctg  640
attcattaga  ggtagaagga  tatacggttg  atgtcaaaaa  680
taaaagaact  tttctttcac  catggatttc  taatattcat  720
gaaaagaaa  gattaaccaa  atataaatca  tctcctgaaa  760
aatggagcac  ggcttctgat  ccgtagactg  atttcgaaaa  800
ggttacagga  cggattgata  agaattgtatc  accagaggca  840
agacaccccc  ttgtggcagc  ttatccgatt  gtacatgtag  880
atatggagaa  tattattctc  tcaaaaaaatg  aggatcaatc  920
cacacagaat  actgatagtg  aaacgagaa  aataagttaa  960
aatacttcta  caagtaggac  acatactagt  gaagtacatg  1000
gaaatgcaga  agtgcattcg  tcgttctttg  atattgggtg  1040
gagtgtatct  gcaggattta  gtaattcgaa  ttcaagtacg  1080
gtcgcatttg  atcattcact  atctctagca  ggggaaagaa  1120
cttgggtcga  aacaatgggt  ttaataatc  ctgatacagc  1160
aagattaaat  gccaatatta  gatattgtaa  tactgggagc  1200
gctccaatct  acaacgtgtt  accaacgact  tcgttagtgt  1240
taggaaaaaa  tcaaacactc  gcgacaatta  aagctaagga  1280
aaaccaatta  agtcaaatat  ttgcacctaa  taattattat  1320
ccttctaaaa  acttggcgcc  aatcgcat  aatgcacaag  1360
acgatttcag  ttctactcca  attacaatga  attacaatca  1400
atttcttgag  ttgaaaaaaa  cgaacaatt  aagatttagat  1440
acggatcaag  tatatgggaa  tatagcaaca  tacaattttg  1480
aaaatggaag  agtgagggtg  gatacaggct  cgaactggag  1520
tgaagtgtta  ccgcaaatc  aagaaacaac  tgcacgtatc  1560
atttttaatg  gaaaagattt  aaatctggta  gaaaggcgga  1600
tagcggcggt  taatcctagt  gatccattag  aaacgactaa  1640
accggatatg  acattaaaag  aagcccttaa  aatagcattt  1680
ggatttaacg  aaccgaatgg  aaacttaca  tatcaaggga  1720
aagacataac  cgaatttgat  ttttaattcg  atcaacaaac  1760
atctcaaaat  atcaagaatc  agttagcgga  attaaacgca  1800
actaacatat  atactgtatt  agataaaatc  aaattaaatg  1840
caaaaaatgaa  tattttaata  agagataaac  gttttcatta  1880
tgatagaaat  aacatagcag  ttggggcgga  tgagttagta  1920
gttaaggagg  ctcatagaga  agtaattaat  tcgtcaacag  1960
agggattatt  gttaaatatt  gataaggata  taagaaaaat  2000
attatcagg  tatattgtag  aaattgaaga  tactgaagg  2040
cttaagaag  ttataaatga  cagatatgat  atgttgaata  2080
tttctagttt  acggcaagat  ggaaaaacat  ttagatatt  2120
taaaaaatat  aatgataaat  taccgttata  tataagtaat  2160
cccaattata  aggtaaatgt  atatgctgtt  actaaagaaa  2200
acactattat  taatcctagt  gagaatgggg  atactgtac  2240
caacgggatc  aagaaaaatt  taatcttttc  taaaaaaggc  2280
tatgagatag  gataa  2295

```

<210> 2

<211> 3011

<212> DNA

<213> *Bacillus anthracis*

<220>

<400> 2

```

atggaagtta aacaggagaa cgggttatta aatgaatcag aatcaagttc
ccaggggtta ctaggatact atttttagtga tttgaatttt caagcaccca
tggtgggttac ctctcttact acaggggat tttctattcc tagttctgag
ttagaaaaata ttccatcgga aaaccaatat ttccaatctg ctatttggtc
aggattttac aaagttaaga agagtgtatga atatacatat gctactccg
tgataaactca tgtaacaaatg tgggtagatg accaagaagt gatataaaa
gctttctaatt ctaacaaaat cagatttagaa aaaggaagat tatatcaaat
aaaaattcaa tatcaacgag aaaatcctac tgaaaaagga ttggtattca
agttgtactg gacggaattct caaaaaataa aagaagtgat tctagtgtat
aacttacaat tgccagaatt aaaaacaaaa tcttcgaact caagaaaaaa
gcgaagtaca agtgctggac ctacgggtcc agacgtgac aatgatggaa
tccttgattc attagaggta gaaggatata cggttgatgt caaaaaataa
agaaacttttc tttcaccatg gatttctaatt attcatgaaa agaaaggatt
aaccaaatat aaatcatctc ctgaaaaatg gagcacggct tctgatccgt
acagtgattt cgaaaaaggtt acaggacgga ttgataagaa tgtatcacca
gaggcaagac acccccttgt ggcagcttat cggattgtac atgtagatat
ggagaatatt attctctcaa aaaaatgagga tcaatccaca cagaatactg
atagtgaaac gagaacaata agtaaaaaata cttctacaag taggcacaat
actagtgaag tacatggaaa tgcagaagtg catgcgtcgt tctttgatat
tggtgggagt gtatctgcag gatttagtaa ttcgaattca agtacggctg
caattgatca ttcactatct ctacgagggg aaagaacttg ggctgaacaa
atgggttttaa ataccgtga tacagcaaga ttaaatgcc aatattagata
tgtaaaatac gggacggctc caatctacaa cgtgttacca acgacttctg
tagtgttagg aaaaaatcaa acactcgcga caattaaagc taaggaaaaa
caattaagtc aaatacttgc acctaataat tattatcctt ctaaaaaact
ggcgccaatc gcattaaatt cacaagacga tttcagttct actccaatta
caatgaatta caatcaattt cttaggttag aaaaaacgaa acaattaaga
ttagatacgg atcaagtata tgggaatata gcaacataca attttgaaaa
tggaagagtg aggggtgata caggctcgaa ctggagtga gttgtaccgc
aaattcaaga aacaactgca cgtatcattt ttaatggaaa agatttaaat
ctggtagaaa ggcggatagc ggcggttaat cctagtgtac cattagaaac
gactaaaccg gatatgacat taaaagaagc cctaaaaata gatttggat
ttaacgaacc gaatggaac ttacaatatc aagggaaaga cataaccgaa
tttgatttta atttcgatca acaaacatct caaaatatca agaatacagt
agcggaatta aacgcaacta acatatatac cgtattagat aaatacaaat
taaatgcaaa aatgaattat ttaataagag ataaacgttt tcattatgat
aggaataaca tagcagttgg ggcggatgag tcagtagtta aggaggtca
tagagaagta attaatcgt caacagaggg attattgtta aatattgata
aggatataag aaaaatatta tcaggttata ttgtagaat tgaagatact
gaagggttta aagaagttat aaatgcaga tatgatattg tgaaatttcc
tagtttacgg caagatggaa aaacatttat agattttaaa aaatatattg
taaatatacc gttatatata agtaatccca aatgtatat ataatgtat
gctgttacta aagaaaacac tattattaat cctagtgaga atggggatac
tagtaccaac gggatcaaga aaattttaat ctttctaaaa aaaggctatg
agataggata a

```

<210> 3

<211> 2292

<212> DNA

<213> *Bacillus anthracis*

<220>

<400> 3

atggatgcaa	tgaagagagg	gctctgctgt	gtgctgctgc	tgtgtgggagc
agtcctcggt	tcggctagcg	aggtgattca	ggcagaagtt	aaacaggaga
accgggttatt	aaatgaatca	gaatcaagtt	cccaggggtt	actaggatatac
tatttttagtg	atttgaatct	tcaagcacc	atgggtggtta	ctctctctac
tacaggggat	ttatctattc	ctagtcttga	gttagaaaaat	attccatcgg
aaaaccaata	ttttcaatct	gctatttggt	caggattttat	caaagttaag
aagagtgtatg	aatatacatt	tgctacttcc	gctgataatc	atgtaacaat
gtgggttagat	gaccaagaag	tgattaataa	agcttctaat	tctaacaaaa
tcagatttaga	aaaaggaaga	ttatatcaaa	taaaaattca	atatcaacga
gaaaaatccta	ctgaaaaagg	attggatttc	aagttgtact	ggaccgattc
tcaaaataaaa	aaagaagtga	tttctagtga	taacttcaaa	ttgccagaat
taaaacaaaa	atcttcgaac	tcaagaaaaa	agcgaagtac	aagtgtctgga
ctcacgggttc	cagaccgtga	caatgatgga	atccctgatt	catttagaggt
agaaggtat	acggttgatg	tcaaaaataa	aagaactttt	ctttaccatt
ggatttctaa	tattcatgaa	aagaaaggat	taaccaataa	taaatcatct
ctgaaaaaat	ggagcaccgc	ttctgtatccg	tacagtgtatt	tcgaaaagggt
tacaggacgg	attgataaga	atgtatcacc	agaggccaaga	accccccttg
tgggcagctta	tcggattgta	catgtagata	tggagaatat	tattctctca
aaaaatgaggt	atcaatccac	acagaatact	gatagtgaaa	cgagaacaat
aagtaaaaaat	acttctacaa	gtaggacaca	tactagttaa	gtacatggaa
atgcagaagt	gcattgcgtcg	ttctttgata	ttgggtgggag	tgtatctgca
ggatttagta	attcgaattc	aagtacggtc	gcaattgatac	attcactatc
tctagcagg	gaaagaactt	gggctgaaac	aatgggttta	aataccgctg
atcacgcaag	attaaatgcc	aatatttagat	atgtaaatac	tgggcagggt
ccaattctaca	acgtgttacc	aacgacttcg	ttagtgttag	gaaaaaatca
aacactcgcg	acaattaaag	ctaaggaaaa	ccaattaaat	caaatcttg
cacctataaa	ttattatcct	tctaaaaact	tggcgccaat	cgcattaaat
gcacaagacg	atttcagttc	tactccaatt	acaatgaatt	acaatcaatt
tccttgagtta	gaaaaaacga	aacaattaaag	attagatagc	gatcaagtat
atgggaatat	agcaacatac	aattttgaaa	atggaagagt	gaggggtgga
acaggctcga	actggagtga	agtggttaccg	caaattcaag	aaacaactgc
acgtatcat	tttaatggaa	aagatttaaa	tctggtagaa	aggcggatag
cggcggttaa	tcctagtgtat	ccattagaaa	cgactaaacc	ggatacgaca
ttaaaagaag	cccttaaaat	agcatttgga	tttaacgaac	cgaatggaaa
cttacaatat	caaggggaaag	acataaccga	atttgatttt	aatttcgcat
aacaacatc	tcaaaatatt	aagaatcagt	tagcggaatt	aaacgcaact
aacatatata	ctgtattaga	taaaatcaaa	ttaaatgcaa	aaatgaatat
tttaataaga	gataaacggt	ttcattatga	tgaaaaatac	atagcagttg
gggcggatga	gtcagtagtt	aaggaggctc	atagagaagt	aattaattcg
tcaacagagg	gattatttgt	aaattatgat	aaggatataa	gaaaaatatt
atcagggttat	attgtagaaa	ttgaagatc	tgaagggtt	aaaggaattt
taaatgcacg	atatgatgat	ttgaatat	ctagtttacg	gcaagatgga
aaaacattta	tagattttta	aaaaataaat	gataaaattac	cgttatatat
aagtaattccc	aattataagg	taaatgtata	tgctgttact	aaagaaaaa

ctattatttaa tcctagtgag aatggggata ctagtaccaa cgggatcaag
 aaaatttttaa tctttttctaa aaaaggctat gagataggat aa

<210> 4

<211> 1710

<212> DNA

<213> *Bacillus anthracis*

<220>

<400> 4

atgagtacaa gtgctggacc tacgggtcca gaccgtgaca atgatggaat
 cctcgattca ttagaggtag aaggatatac ggttgatgtc aaaaaataaa
 gaacttttct ttcaccatgg atttctaata ttcattgaaa gaaaggatta
 accaaatata aatcatctcc tgaaaaatgg agcacggctt ctgatccgta
 cagtgatttc gaaaagggtta caggacggat tgataagaat gtatcaccag
 aggcaagaca cccctctgtg gcagcttatc cgattgtaca tgtagatatg
 gagaatatta ttctctcaaa aatagaggat caatccacac agaatactga
 tagtgaacac agaacaataa gtaaaaaatac ttctacaagt aggacacata
 ctagtgaagt acatggaat gcagaagtgc attgctcgtt ctttgattatt
 ggtgggagtg tatctgcagg atttagtaat tcgaattcaa gtacgggtgc
 aattgatcat tcactatctc tagcagggga aagaacttgg gctgaaacaa
 tgggtttaaa taccgctgat acagcaagat taaatgccaa tattagatat
 gtaaaactcg ggacggctcc aatctacaac gtgttaccaa cgacttcgtt
 agtggttagga aaaaatcaaa cactcgcgac aattaaagct aaggaaaaacc
 aattaagtca aataacttga cctaataatt attatccttc taaaaacttg
 gcgccaatcg cattaaatgc acaagacgat ttcagttcta ctccaattac
 aatgaattac aatcaatttc ttgagttaga aaaaacgaaa caattaaagt
 tagatacggg tcaagtatat gggaaatatag caacatacaa ttttgaatat
 ggaagagtga ggggtggatac aggtcgaac tggagtgaag gtttaccgca
 aattcaagaa acaactgcac gtatcatttt taatggaaaa gatttaaatc
 tggtagaaaag gcggatagcg gcggttaatc ctagtgatcc attagaacacg
 actaaacccg atatgacatt aaaagaagcc cttaaaatag catttgattt
 taacgaaccg aatggaaact tacaatatca agggaaagac ataaccgaat
 ttgattttta ttctgatcaa caaacatctc aaaatatcaa gaatcagtta
 gcggaattaa acgcaactaa catatactat gtattagata aaatcaaat
 aaatgcaaaa atgaatattt taataagaga taaacgtttt cattatgata
 gaaataacat agcagttggg gcggatgagt cagtagttaa ggaggctcat
 agagaagtaa ttaattcgtc aacagaggga ttattgttaa atattgataa
 ggatataaga aaaaatttat caggttatat ttagaaaatt gaagatactg
 aagggcttaa agaagttata aatgacagat atgatattgt gaatatctt
 agtctacggc aagatggaaa aacatttata gatcttaaaa aacataatga
 taaattaccg ttatatataa gtaatcccaa ttataaggta aatgtatatg
 ctgttactaa agaaaacact attattaatc ctagtgaaga tggggatact
 agtaccacac ggatcaagaa aattttaatc ttttctaaaa aaggctatga
 gataggataa

<210> 5

<211> 763

<212> PRT

<213> *Bacillus anthracis*

<220>

<400> 5

1 MDAMKRGLCVLLLCGAVFVSASEVIQAEVKQENRLLNESESS
44 SQGLLGYYFSDLNFQAPMVVTSSTTGDLSIPSSSELENIPSENQYFQSAIWSGFIKVKK
102
SDEYTFATSADNHVTMWVDDQEVINKASNSNKIRLEKGRLYQIKIQYQRENPTTEKGLD
160
FKLYWTDSONKKEVISSDNLQLPELKQKSSNSRKKRSTSAGPTVPDRDNDGIPDSLEV
218
EGYTVDVKNKRTFLSPWISNIHEKKGLTKYKSSPEKWSTASDPYSDFEKVTGRIDKNV
276
SPEARHPLVAAYPIVHVDMENIILSKNEDQSTQNTDSETRTISKNTSTSRHTSEVHG
334
NAEVHASFDDIGGSVSAGFSNSNSSTVAIDHSLSLAGERTWAETMGLNTADTARLNAN
392
IRYVNTGTAPIYNVLPFTSLVLGKNQTLATIKAKENQLSQILAPNNYYPKQNLAPIAL
450
NAQDDFSSTPITMNYNQFLELEKTKQLRLDTDQVYGNIAITYNFENGRVVRVDTGSNWSE
508
VLPQIQETTARIIFNGKDLNLVERRIAAVNPSPDLETTKPDMTLKEALKIAFGFNEPN
566
GNLQYQGGKDITEFDNFQDQTSQNIKNQLAELNATNIYTVLDKIKLNAKMNILIRDKR
624
FHYDRNIAVGADESVVKEAHREVINSSTEGLLLNIDKDIRKILSGYIVEIEDTEGLK
682
EVINDRYDMLNISSLRQDGKTFIDFKKYNDKLPYISPNYKVNVAVTKENTIINPS
740 ENGDTSTNGIKKILIFSCKGYEIG

<210> 6

<211> 764

<212> PRT

<213> *Bacillus anthracis*

<220>

<400> 6

1 MKKKRVLIPLMALSTILVSSTGNLEVIQAEVKQENRLLNESESS

45
SQGLLGYFFSDLNFQAPMVVTSSTTGDLSSIPSELENIPSENQYFQSAIWSGF IKVKK
103
SDEYTFATSADNHVMTMWDDQEVINKASNSNKIRLEKGRLYQIKIQYQRENPTTEKGLD
161
FKLYWTDSONKKEVISSDNLQLPELKQKSSNSRKKRSTSAGPTVPRDRNDGIPDSLEV
219
EGYTVDVKNKRTFLSPWISNIHEKKGLTKYKSSPEKWSTASDPYDFEKTGRIDKNV
277
SPEARHPLVAAYPIVHVDMENIILSKNEDQSTQNTDSETRTISKNTSTSRTHTSEVHG
335
NAEVHASFFDIGGSVSAGFSNSNSSTVAIDHSLSLAGERTWAETMGLNTADTARLNAN
393
IRYVNTGTAPIYNVLPPTSLVLGKNQTLATIKAKENQLSQILAPNNYYPKSNLAPIAL
451
NAQDDFSSTPITMNYNQFLELEKTKQLRLDTDQVYGNIAITYNFENGRVVRVDTGSNWSE
509
VLPQIQETTARIIFNGKDLNLVERRIAAVNPSDPLETTKPDMTLKEALKIAFGNEPN
567
GNLQYQGGKIDITEFDNFDDQTSQNIKNQLAELNATNIYTVLDKIKLNAKNMILIRDKR
625
PHYDRNNIAVGADESUVKEAHREVINSSTEGLLNIDKDIRKILSGYIVEIEDTEGLK
683
EVINDRYDMLNISSLRQDGKTFIDFKKYNDKLPYISNPNYKVNVAVTKENTIINPS
741 ENGDTSTNGIKKILIFSCKGYEIG

<210> 7

<211> 736

<212> PRT

<213> *Bacillus anthracis*

<220>

<400> 7

1 MEVKQENRLNESESS

17
SQGLLGYFFSDLNFQAPMVVTSSTTGDLSSIPSELENIPSENQYFQSAIWSGF IKVKK
75
SDEYTFATSADNHVMTMWDDQEVINKASNSNKIRLEKGRLYQIKIQYQRENPTTEKGLD
133
FKLYWTDSONKKEVISSDNLQLPELKQKSSNSRKKRSTSAGPTVPRDRNDGIPDSLEV
191
EGYTVDVKNKRTFLSPWISNIHEKKGLTKYKSSPEKWSTASDPYDFEKTGRIDKNV
249
SPEARHPLVAAYPIVHVDMENIILSKNEDQSTQNTDSETRTISKNTSTSRTHTSEVHG
307
NAEVHASFFDIGGSVSAGFSNSNSSTVAIDHSLSLAGERTWAETMGLNTADTARLNAN

365
IRYVNTGTAPIYVNLPTTSLVLGKNQTLATIKAKENQLSQILAPNNYPSKNLAPIAL
423
NAQDDFSSTPITMNYNQFLELEKTKQLRLDQVYGNIAATYNFENGVRVDTGSGNWSE
481
VLPQIQETTARIIFNGKDLNLVERRIA AAVNPSPLETTKPDMTLKEALKIAFGFNEPN
539
GNLQYQGDITEFDNFDDQQTSONIKNQLAELNATNIYTVLDKIKLNAKMNILIRDKR
597
FHYDRNNIAVGADES VVKEAHREVINSSTEGLLLNIDKDIRKILSGYIVEIEDTEGLK
655
EVINDRYDMLNIISSLRQDGKTFIDFKKYNDKLPYISNPYKVN VYAVTKENTIINPS
713 ENGDTSTNGIKKILIFS KKGYEIG

<210> 8

<211> 569

<212> PRT

<213> *Bacillus anthracis*

<220>

<400> 8

1 MSTSAGPTVPDRDNDGIPDSLEV
24
EGYTVDVKNKRTFLSPWISNIHEKGLTKYKSSPEKWSTASDPYSDFEKVTGRIDKNV
82
SPEARHPLVAAYPIVHVDMENIILSKNEDQSTQNTDSETRTISKNTSTSRTHTSEVHG
140
NAEVHASFFDIGGSVSAGFSNSNSSTVAIDHSLSLAGERTWAETMGLNTADTARLNAN
198
IRYVNTGTAPIYVNLPTTSLVLGKNQTLATIKAKENQLSQILAPNNYPSKNLAPIAL
256
NAQDDFSSTPITMNYNQFLELEKTKQLRLDQVYGNIAATYNFENGVRVDTGSGNWSE
314
VLPQIQETTARIIFNGKDLNLVERRIA AAVNPSPLETTKPDMTLKEALKIAFGFNEPN
372
GNLQYQGDITEFDNFDDQQTSONIKNQLAELNATNIYTVLDKIKLNAKMNILIRDKR
430
FHYDRNNIAVGADES VVKEAHREVINSSTEGLLLNIDKDIRKILSGYIVEIEDTEGLK
488
EVINDRYDMLNIISSLRQDGKTFIDFKKYNDKLPYISNPYKVN VYAVTKENTIINPS
546 ENGDTSTNGIKKILIFS KKGYEIG

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 99/15568

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C12N15/31 C07K14/32 A61K39/07 C12N7/01 C12N1/21

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 A61K C07K C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	L.C. IACONO-CONNORS ET AL.: "PROTECTION AGAINST ANTHRAX WITH RECOMBINANT VIRUS-EXPRESSED PROTECTIVE ANTIGEN IN EXPERIMENTAL ANIMALS" INFECTION AND IMMUNITY, vol. 59, no. 6, June 1991 (1991-06), pages 1961-1965, XP000916157	1, 2, 4
Y	the whole document --- -/--	4-20

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

24 July 2000

Date of mailing of the international search report

02/08/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Julia, P

INTERNATIONAL SEARCH REPORT

Inter nal Application No
PCT/US 99/15568

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	PUSHKO P ET AL: "Replicon-helper systems from attenuated Venezuelan Equine Encephalitis virus: Expression of heterologous genes in vitro and immunization against heterologous pathogens in vivo" VIROLOGY,US,ACADEMIC PRESS,ORLANDO, vol. 239, no. 2, 22 December 1997 (1997-12-22), pages 389-401-401, XP002123951 ISSN: 0042-6822 the whole document ---	4-20
A	WO 96 17067 A (ZYMOGENETICS INC) 6 June 1996 (1996-06-06) page 14, line 18 -page 16, line 35 ---	1-5, 7, 10-20
X	SINGH Y ET AL: "A DELETED VARIANT OF BACILLUS ANTHRACIS PROTECTIVE ANTIGEN IS NON- TOXIC AND BLOCKS ANTHRAX TOXIN ACTION IN VIVO" JOURNAL OF BIOLOGICAL CHEMISTRY,US,AMERICAN SOCIETY OF BIOLOGICAL CHEMISTS, BALTIMORE, MD, vol. 264, no. 32, 15 November 1989 (1989-11-15), pages 19103-19107, XP000650160 ISSN: 0021-9258 the whole document ---	1-3
Y	the whole document	4-20
Y	WO 95 07994 A (VIAGENE INC) 23 March 1995 (1995-03-23) page 9 lines 10-12, page 33 lines 31-35 and example 6 ---	4-20
X	IVINS B E ET AL: "CLONING AND EXPRESSION OF THE BACILLUS ANTHRACIS PROTECTIVE ANTIGEN IN BACILLUS SUBTILIS" INFECTION AND IMMUNITY,US,AMERICAN SOCIETY FOR MICROBIOLOGY, WASHINGTON, vol. 54, no. 2, 1 November 1986 (1986-11-01), pages 537-542, XP000650052 ISSN: 0019-9567 the whole document ---	1-3
Y	the whole document	4-20
Y	WO 96 37616 A (UNIV NORTH CAROLINA ;US HEALTH (US); JOHNSTON ROBERT E (US); DAVIS) 28 November 1996 (1996-11-28) the whole document ---	4-20

	-/-	

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 99/15568

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 98 08952 A (SECR DEFENCE ;BAILLIE LESLIE WILLIAM JAMES (GB)) 5 March 1998 (1998-03-05) the whole document -----</p>	1-3

INTERNATIONAL SEARCH REPORT

Information on patent family members

Inter- national Application No
PCT/US 99/15568

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9617067 A	06-06-1996	US 5641655 A	24-06-1997
		AU 689373 B	26-03-1998
		AU 4737696 A	19-06-1996
		CA 2206399 A	06-06-1996
		CN 1171819 A	28-01-1998
		EP 0799317 A	08-10-1997
		NZ 301858 A	28-01-1999
WO 9507994 A	23-03-1995	AU 3682699 A	07-10-1999
		AU 703653 B	01-04-1999
		AU 5645098 A	04-06-1998
		AU 690583 B	30-04-1998
		AU 7835894 A	03-04-1995
		CA 2158937 A	23-03-1995
		EP 0694070 A	31-01-1996
		EP 0711829 A	15-05-1996
		EP 0716148 A	12-06-1996
		EP 0814154 A	29-12-1997
		EP 0982405 A	01-03-2000
		FI 954601 A	23-02-1996
		JP 9503657 T	15-04-1997
		NO 953901 A	04-01-1996
		US 6015686 A	18-01-2000
		US 5814482 A	29-09-1998
		US 5843723 A	01-12-1998
		US 5789245 A	04-08-1998
		US 6015694 A	18-01-2000
WO 9637616 A	28-11-1996	US 5792462 A	11-08-1998
		AU 5925696 A	11-12-1996
		CA 2220964 A	28-11-1996
		EP 1012295 A	28-06-2000
		JP 11505719 T	25-05-1999
WO 9808952 A	05-03-1998	EP 0927255 A	07-07-1999
		GB 2331523 A	26-05-1999